

UNCLASSIFIED

AD 401 222

*Reproduced
by the*

DEFENSE DOCUMENTATION CENTER

FOR

SCIENTIFIC AND TECHNICAL INFORMATION

CAMERON STATION, ALEXANDRIA, VIRGINIA



UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

63-3-2

MEMORANDUM

RM-3570-PR


APR 11 1963

CATALOGED BY ASTIA
AS AD NO. 401222

401 222

A UNIQUENESS CONDITION FOR
NONTRIVIAL PERIODIC SOLUTIONS TO
THE LIÉNARD EQUATION

T. A. Brown



PREPARED FOR:

UNITED STATES AIR FORCE PROJECT RAND

The **RAND** Corporation
SANTA MONICA • CALIFORNIA

MEMORANDUM

RM-3570-PR

APRIL 1963

**A UNIQUENESS CONDITION FOR
NONTRIVIAL PERIODIC SOLUTIONS TO
THE LIÉNARD EQUATION**

T. A. Brown

This research is sponsored by the United States Air Force under Project RAND—contract No. AF 49(638)-700 monitored by the Directorate of Development Planning, Deputy Chief of Staff, Research and Development, Hq USAF. Views or conclusions contained in this Memorandum should not be interpreted as representing the official opinion or policy of the United States Air Force.

The **RAND** *Corporation*

1700 MAIN ST. • SANTA MONICA • CALIFORNIA

PREFACE

Part of the Project RAND research program consists of basic supporting studies in mathematics. The mathematical research presented here concerns the periodic solutions of the Liénard differential equation, which applies to a wide variety of physical problems.

SUMMARY

This Memorandum presents a new condition which implies that the differential equation

$$\ddot{x} + f(x)\dot{x} + q(x)x = 0$$

has an essentially unique non-trivial periodic solution, to which all other solutions tend as $t \rightarrow \infty$.

ACKNOWLEDGMENT

The author wishes to acknowledge his appreciation to G. Birkhoff and N. Levinson for their help and encouragement in this research.

A UNIQUENESS CONDITION FOR NONTRIVIAL PERIODIC SOLUTIONS TO THE LIÉNARD EQUATION

The existence and uniqueness of periodic solutions to the equation

$$(1) \quad \ddot{x} + f(x)\dot{x} + q(x)x = 0 \quad (f(x), q(x) \text{ continuous})$$

have been widely discussed during the past thirty-five years, notably by Liénard [5] and by Levinson and Smith [4]. The purpose of the present paper is to present a uniqueness condition for periodic solutions which includes one of those given in the latter paper [4] as a special case.

Define the following functions:

$$F(x) = \int_0^x f(x)dx,$$

$$Q(x) = \int_0^x q(x)xdx.$$

In the Poincaré phase-plane (i.e., the (x, \dot{x}) -plane), define (as in [4]) a pseudo-energy function

$$E(x, \dot{x}) = \frac{1}{2}(\dot{x} + F(x))^2 + Q(x).$$

Using this energy function, the following result may be proved:

Lemma. Suppose the following conditions hold on the coefficients of equation (1):

- (a) $q(x) > 0$ for all $x \neq 0$;
- (b) there exist real numbers $a < 0$ and $b > 0$
such that $F(a) = F(b) = F(0) = 0$, and
 $xF(x) < 0$ for all other x such that $a \leq x \leq b$;
- (c) $f(x) \geq 0$ for $x < a$ and $x > b$;
- (d) $\lim_{x \rightarrow \infty} Q(x) = \lim_{x \rightarrow -\infty} Q(-x) = \infty$.

Then there exists at least one nontrivial periodic solution, and at most one which enters both the region
 $x \leq a$ and the region $x \geq b$.

Proof. Along solution curves, $dE/dt = -xq(x)F(x)$. Thus the phase-plane is divided into three strips (see Fig. 1): $x < a$ (where $dE/dt < 0$), $a \leq x \leq b$ (where $dE/dt \geq 0$), and $x > b$ (where $dE/dt < 0$). Now let Γ_0 be a limit cycle which enters all three regions, and suppose it passes through the point $(0, \dot{x}_0)$. Its journey around the phase-plane naturally divides into five parts (as in the figure).

Now say $x_A > x_0$. Let Γ_A be the curve through $(0, x_A)$. In part I of Γ_A 's journey about the phase-plane, it is gaining energy. Since, however, $\dot{x}_{\Gamma_A} > \dot{x}_{\Gamma_0}$ for corresponding values of x in part I, it follows that the change in E per unit x is less on Γ_A than on Γ_0 (for corresponding x). Thus the amount of energy gained by




Fig. 1 — The proof of the lemma

going from $x = 0$ to $x = b$ along Γ_A is less than that gained by going along Γ_0 . Similarly, Γ_A gains less energy than Γ_0 in parts III and V. A simple computation ([1], [2]) shows that Γ_A loses more energy than Γ_0 in parts II and IV. Thus Γ_A must, on balance, lose energy in making a single circuit.

Similarly any Γ_B which starts inside Γ_0 will gain energy in making a circuit provided it enters both the left-hand and right-hand regions. Thus, if such a limit cycle exists, it is unique and stable on both sides.

To show a limit cycle exists, we need only find an \dot{x}_B so small that the cycle starting at $(0, \dot{x}_B)$ gains energy, and an \dot{x}_A so large that the cycle starting at $(0, \dot{x}_A)$ loses energy. This is easy. Note, however, that such a limit cycle need not enter all three regions (as we assumed Γ_0 did [1], [3]), and thus we cannot conclude that there is only one limit cycle unless we insure that any limit cycle will enter both the region $x < a$ and the region $x > b$. Levinson and Smith met this problem by assuming $f(x)$ and $q(x)$ symmetric about zero, but we shall show that a much weaker condition is adequate.

Theorem 1. Suppose an equation satisfying the conditions of the lemma satisfies also the following:

$$(*) \int_0^a q(x)x \, dx = \int_0^b q(x)x \, dx.$$

Then there exists a unique (up to translations in t) non-trivial periodic solution, to which all other solutions tend.

Proof. We have already shown that there is at most one limit cycle which enters both the regions $x < a$ and $x > b$. We have also shown that some limit cycle exists. Now we will show that under condition (*), any limit cycle enters both the above regions, and thus uniqueness follows. First note (see Fig. 2) that along the curve $v = -F(x)$,

$$E(x, v) = \int_0^x q(x)x \, dx.$$

Since $\int_0^x q(x)x \, dx$ is monotone decreasing for $x < 0$, and monotone increasing for $x > 0$, it follows that the energy along the curve $v = -F(x)$, $a < x < b$, is always less than

$$\int_0^a q(x)x \, dx = \int_0^b q(x)x \, dx.$$

Now, consider the orbit I (which starts at $(a, 0)$) or the orbit II (which starts at $(b, 0)$). The energy E is increasing along these orbits, and thus neither of them can intersect $v = -F(x)$ between a and b , since each orbit has initial energy $\int_0^a q(x)x \, dx$. Thus, I must enter $x > b$ and II must enter $x < a$. It immediately follows that any limit cycle must do likewise, and the proof is complete.




Fig. 2—The proof of the theorem

Corollary. Given an equation of the form (1) which satisfies the following conditions:

- (a) $f(x) = f(-x)$, $q(x) = q(-x)$ for all x ;
- (b) $q(x)x$ is differentiable for all x , and $q(x) > 0$ for $x \neq 0$;
- (c) there exists an $a > 0$ such that $f(x) > 0$ for $x > a$;
- (d) $F(x) < 0$ for $0 < x < a$, $F(a) = 0$;
- (e) $\lim_{x \rightarrow \infty} F(x) = \lim_{x \rightarrow \infty} Q(x) = +\infty$.

Then there exists a unique (up to translations in t) non-trivial periodic solution.

Proof. Assumption (a) shows that $F(a) = F(-a)$ and $Q(a) = Q(-a)$, and thus the corollary is easily seen to follow. This corollary is the theorem (unnumbered) which Levinson and Smith prove in Section 4 of [4].

We conclude by giving a specific example of an equation which has a unique stable limit cycle by our theorem, but about which the Levinson-Smith theorems are silent:

$$\ddot{x} + (3x^2 - 4x - 3)\dot{x} + q(x)x = 0$$

$$\text{where } q(x) = \begin{cases} -27x & \text{for } x \leq 0 \\ x & \text{for } x > 0 \end{cases}$$

(here $a = -1$, $b = 3$).

REFERENCES

1. Brown, T. A., "The Asymptotic Behavior of Some Non-linear Autonomous Systems," unpublished doctoral thesis, Harvard University, 1962.
2. Coddington, E. A., and N. Levinson, Theory of Ordinary Differential Equations, New York, 1955.
3. de Figueiredo, R. P., Existence and uniqueness of the periodic solution of an equation for autonomous oscillations, Contributions to the Theory of Non-linear Oscillations, Vol. 5, Princeton University Press, Princeton, N. J., 1960, pp. 269-284.
4. Levinson, N., and O. K. Smith, "A General Equation for Relaxation Oscillations," Duke Math. Jour., Vol. 9, 1942, pp. 382-405.
5. Lienard, A., "Etude des oscillations entretenues," Rev. Gen. d'Elect., Vol. 28, 1928, pp. 901-946.